

THE ANTIPODE OF A DUAL QUASI-HOPF ALGEBRA WITH NONZERO INTEGRALS IS BIJECTIVE

M. BEATTIE, M.C. IOVANOV, AND Ş. RAIANU

Dedicated to Fred Van Oystaeyen for his sixtieth birthday

ABSTRACT. For A a Hopf algebra of arbitrary dimension over a field K , it is well-known that if A has nonzero integrals, or, in other words, if the coalgebra A is co-Frobenius, then the space of integrals is one-dimensional and the antipode of A is bijective. Bulacu and Caenepeel recently showed that if H is a dual quasi-Hopf algebra with nonzero integrals, then the space of integrals is one-dimensional, and the antipode is injective. In this short note we show that the antipode is bijective.

1. INTRODUCTION

The definition of quasi-Hopf algebras and the dual notion of dual quasi-Hopf algebras is motivated by quantum physics and dates back to work of Drinfel'd [4]. The theory of integrals for quasi-Hopf algebras was studied in [9, 6, 2]. In [2], Bulacu and Caenepeel showed that a dual quasi-Hopf algebra is co-Frobenius as a coalgebra if and only if it has a nonzero integral. In this case, the space of integrals is one-dimensional and the antipode is injective, so that for finite dimensional dual quasi-Hopf algebras the antipode is bijective. In this note, we use the ideas from a new short proof of the bijectivity of the antipode for Hopf algebras by the second author [7] to show that the antipode of a dual quasi-Hopf algebra with integrals is bijective, thus extending the classical result of Radford [10] for Hopf algebras.

In this paper we prove

Theorem 1.1. *Let H be a co-Frobenius dual quasi-Hopf algebra, equivalently, a dual quasi-Hopf algebra having nonzero integrals. Then the antipode of H is bijective.*

2. PRELIMINARIES

In this section we briefly review the definition of a dual quasi-Hopf algebra over a field K . We refer the reader to [1, 3, 11] for the basic definitions and properties of coalgebras and their comodules and of Hopf algebras. For the definition of dual quasi-Hopf algebra we follow [8, Section 2.4].

Definition 2.1. *A dual quasi-bialgebra H over K is a coassociative coalgebra (H, Δ, ε) together with a unit $u : K \rightarrow H$, $u(1) = 1$, and a not necessarily associative multiplication $M : H \otimes H \rightarrow H$. The maps u and M are coalgebra maps. We write ab for $M(a \otimes b)$.*

The first author's research was supported by an NSERC Discovery Grant.

The second author was partially supported by the contract nr. 24/28.09.07 with UEFISCU "Groups, quantum groups, corings and representation theory" of CNCIS, PN II (ID_1002).

As well, there is an element $\varphi \in (H \otimes H \otimes H)^*$ called the reassociator, which is invertible with respect to the convolution algebra structure of $(H \otimes H \otimes H)^*$. The following relations must hold for all $h, g, f, e \in H$:

$$\begin{aligned} (1) \quad & h_1(g_1 f_1) \varphi(h_2, g_2, f_2) = \varphi(h_1, g_1, f_1)(h_2 g_2) f_2 \\ (2) \quad & 1h = h1 = h \\ (3) \quad & \varphi(h_1, g_1, f_1 e_1) \varphi(h_2 g_2, f_2, e_2) = \varphi(g_1, f_1, e_1) \varphi(h_1, g_2 f_2, e_2) \varphi(h_2, g_3, f_3) \\ (4) \quad & \varphi(h, 1, g) = \varepsilon(h) \varepsilon(g) \end{aligned}$$

Here we use Sweedler's sigma notation with the summation symbol omitted.

Definition 2.2. A dual quasi-bialgebra H is called a dual quasi-Hopf algebra if there exists an antimorphism S of the coalgebra H and elements $\alpha, \beta \in H^*$ such that for all $h \in H$:

$$\begin{aligned} (5) \quad & S(h_1) \alpha(h_2) h_3 = \alpha(h) 1, \quad h_1 \beta(h_2) S(h_3) = \beta(h) 1 \\ (6) \quad & \varphi(h_1 \beta(h_2), S(h_3), \alpha(h_4) h_5) = \varphi^{-1}(S(h_1), \alpha(h_2) h_3, \beta(h_4) S(h_5)) = \varepsilon(h). \end{aligned}$$

Let H be a dual quasi-Hopf algebra. As in the Hopf algebra case, a left integral on H is an element $T \in H^*$ such that $h^* T = h^*(1) T$ for all $h^* \in H^*$; the space of left integrals is denoted by $\int_l$ and by [2, Proposition 4.7] has dimension 0 or 1. Right integrals are defined analogously with space of right integrals denoted by $\int_r$. Suppose $0 \neq T \in \int_l$. It is easily seen that $\int_l$ is a two sided ideal of the algebra H^* , and $KT \subseteq \text{Rat}(H^*)$ with right comultiplication given by $T \mapsto T \otimes 1$. Since for co-Frobenius coalgebras $\text{Rat}(H^*) = \text{Rat}(H^* H^*) = \text{Rat}(H_{H^*}^*)$, KT must have left comultiplication $T \mapsto a \otimes T$. By coassociativity, a is a grouplike element, called the distinguished grouplike of H . Then, for all $h^* \in H^*$,

$$(7) \quad Th^* = h^*(a)T.$$

From [2, Proposition 4.2], the function $\theta^* : \int_l \otimes H \rightarrow \text{Rat}(H^* H^*)$

$$(8) \quad \theta^*(T \otimes h) = \sigma(S(h_5) \otimes \alpha(h_6) h_7) (S(h_4) \rightharpoonup T) \sigma^{-1}(S(h_3) \otimes \beta(S(h_2)) S^2(h_1))$$

is an isomorphism of right H -comodules, where $\sigma : H \otimes H \rightarrow H^*$ is defined by $\sigma(h \otimes g)(f) = \varphi(f, h, g)$, σ^{-1} is the convolution inverse of σ , and, as usual, $(h \rightharpoonup T)(g) = T(gh)$.

3. PROOF OF THE THEOREM

Let H be a quasi-Hopf algebra with $0 \neq T \in \int_l$. As in [7], for each right H -comodule (M, ρ) , we denote by ${}^a M$ the left H -comodule structure on M defined by $m_{-1}^a \otimes m_0^a = aS(m_1) \otimes m_0$, where $\rho(m) = m_0 \otimes m_1$. Denote the induced right H^* -module structure on ${}^a M$ by $m \cdot^a h^* = h^*(m_{-1}^a) m_0^a = h^*(aS(m_1)) m_0$. By [2, Corollary 4.4] the antipode S of H is injective, and therefore has a left inverse S^l . Then, for σ as above, we have the following analogue of [7, Proposition 2.5]:

Proposition 3.1. *The map $p : {}^a H \rightarrow \text{Rat}(H^*)$ defined by*

$$(9) \quad p(h) = \sigma(S(S^l(h_3)) \otimes \alpha(S^l(h_2))S^l(h_1)) * (h_4 \rightharpoonup T) * \sigma^{-1}(h_5\beta(h_6) \otimes S(h_7))$$

is a surjective morphism of left H -comodules.

Proof. Let $\Psi := \sigma(S(S^l(h_3)) \otimes \alpha(S^l(h_2))S^l(h_1))$. Then for $c^* \in H^*$ and $g \in H$:

$$\begin{aligned} (p(h) * c^*)(g) &= p(h)(g_1)c^*(g_2) \\ (9) &= \Psi(g_1)T(g_2h_4)\sigma^{-1}(h_5\beta(h_6) \otimes S(h_7))(g_3)c^*(g_4) \\ &= \Psi(g_1)T(g_2h_4)\varphi^{-1}(g_3, h_5\beta(h_6), S(h_7))c^*(g_4) \\ &= \Psi(g_1)T(g_2h_4)\varphi^{-1}(g_3, h_5, S(h_7))c^*(g_4\beta(h_6)) \\ (5) &= \Psi(g_1)T(g_2h_4)\varphi^{-1}(g_3, h_5, S(h_9))c^*(g_4(h_6\beta(h_7)S(h_8))) \\ &= \Psi(g_1)T(g_2h_4)c^*(\varphi^{-1}(g_3, h_5, S(h_9))g_4(h_6\beta(h_7)S(h_8))) \\ &= \Psi(g_1)T(g_2h_4)\beta(h_7)c^*(\varphi^{-1}(g_3, h_5, S(h_9))g_4(h_6S(h_8))) \\ (1) &= \Psi(g_1)T(g_2h_4)\beta(h_7)c^*((g_3h_5)S(h_9)\varphi^{-1}(g_4, h_6, S(h_8))) \\ &= \Psi(g_1)T(g_2h_4)(S(h_9) \rightharpoonup c^*)(g_3h_5)\beta(h_7)\varphi^{-1}(g_4, h_6, S(h_8)) \\ &= \Psi(g_1)(T * (S(h_8) \rightharpoonup c^*))(g_2h_4)\beta(h_6)\varphi^{-1}(g_3, h_5, S(h_7)) \\ (7) &= \Psi(g_1)(S(h_8) \rightharpoonup c^*)(a)T(g_2h_4)\beta(h_6)\varphi^{-1}(g_3, h_5, S(h_7)) \\ &= \Psi(g_1)T(g_2h_4)\sigma^{-1}(h_5\beta(h_6) \otimes S(h_7))(g_3)c^*(aS(h_8)) \\ (9) &= p(h_1)(g)c^*(aS(h_2)) \\ &= p(c^*(aS(h_2))h_1)(g) \\ &= p(c^*(h_{-1}^a)h_0^a)(g) \\ &= p(h \cdot^a c^*)(g). \end{aligned}$$

Thus p is left H -colinear. Finally, we note that $p \circ S = \theta^*(T \otimes -)$ where θ^* is the isomorphism from (8) so that p is surjective. $\square$

Let c be a grouplike element of H . From [2, p.580], c is invertible with inverse $S(c)$. We will show that left multiplication by c has an inverse too.

Let $\theta_c \in \text{End}(H)$ be defined by $\theta_c(h) = ch$ and define the coinver automorphisms q_c and $r_c = q_c^{-1} \in \text{End}(H)$ by:

$$q_c(h) = \varphi^{-1}(c, S(c), h_1)h_2\varphi(c, S(c), h_3) \text{ and } r_c(h) = \varphi(c, S(c), h_1)h_2\varphi^{-1}(c, S(c), h_3).$$

Lemma 3.2. *For any grouplike element c and θ_c, r_c, q_c as above, $\theta_c \circ \theta_{c^{-1}} = r_c$ and thus θ_c is bijective with inverse $\theta_c^{-1} = \theta_{c^{-1}} \circ q_c = q_{c^{-1}} \circ \theta_{c^{-1}}$.*

Proof. Using (1) and the fact that $c^{-1} = S(c)$, we see that

$$\theta_c \circ \theta_{c^{-1}}(h) = c(c^{-1}h) = \varphi(c, S(c), h_1)(cS(c))h_2\varphi^{-1}(c, S(c), h_3) = r_c(h).$$

The same formula for $c^{-1} = S(c)$ yields $\theta_{c^{-1}} \circ \theta_c = r_{c^{-1}}$ and the statement then follows directly. $\square$

We can now prove our main result.

Proof of Theorem 1.1.

We only need to prove the surjectivity. The proof goes along the lines of the proof of [7, Theorem 2.6], but with the difference that here the antipode is not necessarily an anti-morphism of algebras.

Let π be the composition map ${}^aH \xrightarrow{p} \text{Rat}(H_{H*}^*) \xrightarrow{\sim} H \otimes \int_r \simeq H$, where the last two isomorphisms follow by left-right symmetry of the results of [2]. Since H is a co-Frobenius coalgebra, ${}^H H$ is projective by [5, Theorem 1.3] or [2, Theorem 4.5, (x)], and as π is surjective, there is a morphism of left H -comodules $\lambda : H \rightarrow {}^aH$ such that $\pi\lambda = \text{Id}_H$. We then have

$$aS(\lambda(h)_2) \otimes \lambda(h)_1 = \lambda(h)_{-1}^a \otimes \lambda(h)_0^a = h_1 \otimes \lambda(h_2).$$

Applying $\text{Id} \otimes \varepsilon\pi$, we get $aS(\varepsilon\pi(\lambda(h)_1)\lambda(h)_2) = h$ for any $h \in H$. Thus $\theta_a \circ S$ is surjective and since θ_a is bijective by Lemma 3.2, S is surjective also. $\square$

REFERENCES

- [1] E. Abe, *Hopf Algebras*, Cambridge Univ. Press, 1977.
- [2] D. Bulacu, S. Caenepeel, Integrals for (dual) quasi-Hopf algebras. Applications, J. Algebra **266** (2003), no. 2, 552-583.
- [3] S. Dăscălescu, C. Năstăsescu, Ș. Raianu, *Hopf Algebras: an Introduction*, Monographs and Textbooks in Pure and Applied Mathematics **235**, Marcel Dekker, Inc., New York, 2001.
- [4] V.G. Drinfel'd, Quasi-Hopf Algebras, Leningrad Math. J. **1** (1990) 1419-1457.
- [5] J. Gomez-Torrecillas, C. Năstăsescu, Quasi-co-Frobenius coalgebras, J. Algebra **174** (1995), no. 3, 909-923.
- [6] F. Hausser, F. Nill, Integral theory for quasi-Hopf algebras, arXiv:math/9904164v2.
- [7] M.C. Iovanov, Generalized Frobenius algebras and the theory of Hopf algebras, preprint, arXiv:0803.0775.
- [8] S. Majid, *Foundations of Quantum Group Theory*, Cambridge University Press, Cambridge, 1995.
- [9] F. Panaite, F. van Oystaeyen, Existence of integrals for finite dimensional Hopf algebras, Bull. Belg. Math. Soc. Simon Stevin **7** (2000) 261-264.
- [10] D.E. Radford, Finiteness conditions for a Hopf algebra with a nonzero integral, J. Algebra **46** (1977), no. 1, 189-195.
- [11] M.E. Sweedler, *Hopf Algebras*, Benjamin, New York, 1969.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, MOUNT ALLISON UNIVERSITY,
SACKVILLE, NEW BRUNSWICK, CANADA E4L 1E6
E-mail address: mbeattie@mta.ca

STATE UNIVERSITY OF NEW YORK, 244 MATHEMATICS BUILDING, BUFFALO, NY 14260-2900,
USA, AND UNIVERSITY OF BUCHAREST, FAC. MATEMATICA & INFORMATICA, STR. ACADEMIEI
NR. 14, BUCHAREST 010014, ROMANIA
E-mail address: yovanov@gmail.com

MATHEMATICS DEPARTMENT, CALIFORNIA STATE UNIVERSITY, DOMINGUEZ HILLS, 1000 E VIC-
TORIA ST, CARSON CA 90747, USA
E-mail address: sraianu@csudh.edu